

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/280641159>

CHRONIC TOXICITY OF CYPERMETHRIN, A PYRETHROID INSECTICIDE TO EPIGEIC EARTHWORM EISENIA FETIDA

Article in *International Journal of Current Research* · August 2015

CITATION

1

READS

445

1 author:



Partha Pratim Chakravorty

Raja Narendralal Khan Women's College(Autonomous)

70 PUBLICATIONS 221 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



1. Ecophysiological--Soil microarthropod . [View project](#)



Role of aerobic bacteria in the gut of selected species of earthworm in the degradation of pesticides in soil [View project](#)



ISSN: 0975-833X

RESEARCH ARTICLE

CHRONIC TOXICITY OF CYPERMETHRIN, A PYRETHROID INSECTICIDE TO EPIGEIC
EARTHWORM *EISENIA FETIDA*

Rupa Dasgupta, Somanka Sanyal and *Partha P. Chakravorty

Department of Zoology, Raja N. L. Khan Women's College, Midnapur, Paschim Medinipore -721102,
West Bengal, India

ARTICLE INFO

Article History:

Received 22nd April, 2015
Received in revised form
05th May, 2015
Accepted 19th June, 2015
Published online 28th July, 2015

Key words:

Sub-lethal dose;
Eisenia fetida;
Biomass; Cocoon;
Juveniles;
Alkaline phosphatase.

ABSTRACT

Effects of sub-lethal doses of cypermethrin on biomass, reproduction and alkaline phosphatase activity of the epigeic earthworm, *Eisenia fetida* were investigated in natural grassland soil (pH-6.72±0.02, organic carbon-1.28±0.37) under laboratory conditions. *E. fetida* exposed to 0.006 to 0.025 mg/kg soil of cypermethrin, corresponding to 12.5 (T2) to 50 % (T4) of LC₅₀ values of cypermethrin did not exhibit any significant reduction in biomass from control. Production of cocoons and juveniles remained significantly unaltered at T2 and T3 doses of the insecticide but was inhibited with exposure to the highest dose (T4) of cypermethrin. Alkaline phosphatase activity was found to remain significantly unaltered at T2 but increased both in T3 and T4 in comparison to control. Thus, though the LC₅₀ value of cypermethrin was much higher than its recommended agricultural dose indicating its less hazardous nature, it can be concluded from the present study that sub-lethal doses of cypermethrin significantly affected reproduction and alkaline phosphatase activity of the test specimen and their widespread application in agro-ecosystems must be carefully monitored. The present study also indicated the potential use of alkaline phosphatase as a biomarker in assessment of insecticide pollution.

Copyright © 2015 Rupa Dasgupta et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Rupa Dasgupta, Somanka Sanyal and Partha P. Chakravorty, 2015. "Chronic toxicity of cypermethrin, a pyrethroid insecticide to epigeic earthworm *Eisenia fetida*", *International Journal of Current Research*, 7, (7), 17693-17696.

INTRODUCTION

The urge for increasing agricultural yield to feed the increasing population has brought a large number of chemicals in protecting the crop against insect pests. However, a grave environmental problem have aroused because of the indiscriminate usage of these chemicals in agricultural fields (Tillman and Mulrooney, 2000; Zhu et al., 2004; Reinecke and Reinecke, 2007). Insecticides are known to be toxic to many non-target organisms and also cause serious sub-lethal effects (Nath et al., 1997; Suh et al., 2000; Yasmin, 2007) including increases and decreases in reproductive potential and growth rate (Takada et al., 2001; Willrich and Boethel, 2001; Yasmin and Dsouza, 2010) Cypermethrin is a pyrethroid insecticide that is used to control insect pests (Cox, 1996) and kills insects by disrupting normal functioning of the nervous system (Vijenbergh & Bercken, 1990). Though it has been shown that they are harmful to beneficial insects, pyrethroids are effective

at low rates and relatively inexpensive (Nowak et al., 2001; Xu et al., 2001). However the effects of pyrethroids on earthworms, a major group of non-target soil fauna has not been fully explored. Cypermethrin, one of the major pyrethroid insecticides currently used worldwide against insect pests (Usmani and Knoewles, 2001). Has been used in the present study to determine its effects on a non-target soil fauna *Eisenia fetida*, an epigeic earthworm known for its efficacy in nutrient recycling and widely used for vermicompost production throughout the world

MATERIALS AND METHODS

Specimens of *Eisenia fetida* were collected from a local vermicompost unit around Midnapore town (W. Bengal, India) and were brought to the laboratory in plastic bags along with compost materials, identified and only pure species of *E. fetida* were used for culture. Specimens were cultured in large earthen pots (diameter 45.72 cm, mean depth 30.48 cm) with finely ground soil (collected from grassland that has never been used for any agricultural purpose and pest control; pH - 6.72 ± 0.02, organic carbon-1.28±0.37) and farmyard manure mixed in the ratio of 1:1 was used as the culture medium. The

*Corresponding author: Partha P. Chakravorty
Department of Zoology, Raja N. L. Khan Women's College, Midnapur,
Paschim Medinipore -721102, West Bengal, India

culture pots were covered with fine meshed iron nets and kept inside BOD incubators at $28 \pm 0.5^\circ \text{C}$. An approximate level of 60-70 % moisture was maintained by adding distilled water into the medium. Finely ground farmyard manure was added as feed every week during the entire period of culture. The cocoons were hand sorted, cultured in separate culture pots and were later used as test specimens following the guidelines for testing of chemicals for earthworm (OECD, 1984; 2004). The pH of the soil was measured using glass electrode pH meter (Systronics model 512SE) and organic carbon content was determined following the methods of (Walkley and Black, 1934). Cypermethrin ((RS)-a cyano-3phenoxy benzyl(1RS)-cis, trans-3 (2,2-dichlorovinyl)- 2,2- dimethylcyclopropane carboxylate) a synthetic derivative of Pyrethrum, a stomach and contact insecticide effective against a broad range of pests of cotton, fruit and vegetable crops was used as the test chemical. Experiments were conducted in small inert polythene boxes (total area, 192 cm^2) containing 600 g of finely ground grassland soil previously used in the culture medium as the test medium for the toxicity studies. Sub lethal doses of the insecticides were applied based on the previously established LC_{50} value of the insecticide on *E. fetida* (Dasgupta et al., 2011). The experimental boxes were exposed to control (T1) and different doses of insecticides (T2 to T4) representing 12.5, 25 and 50 % LC_{50} values of cypermethrin in five replicates (Table 1).

Sublethal doses of pesticides

Table 1. Doses (mg formulation /kg soil) of cypermethrin used in the sub lethal toxicity

Dose	Cypermethrin (mg/kg soil)
T ₁ (control)	0.000
T ₂	0.006
T ₃	0.012
T ₄	0.025

Age-synchronized adult specimens of *E. fetida* (average wet weight 300-350 mg) were collected and 10 such specimens were placed in each experimental box. Before introduction, the worms were rinsed with water, blotted dry on a filter paper and biomass per ten worms was determined. The boxes were maintained at $28 \pm 0.5^\circ \text{C}$ inside a BOD incubator and 5 g (dry weight) of finely ground cow manure moistened to 50 % w/w was added each week to provide food for the growing worms. Additional food was given when all the food added was consumed. Moisture loss from the test soil was checked by weighing the test containers at weekly intervals and replenished if needed. The worms were weighed on the 28th day to determine the change in biomass and were removed from the test boxes. Reproductive success of the test specimens was determined from the rate of cocoon and juvenile production. For determination of rate of cocoon production the test medium was carefully examined with a magnifying glass every week and number of cocoons was noted. The boxes were left undisturbed after removal of the adults on the 28th day till the 56th day and were housed inside a BOD incubator. The temperature and moisture were maintained at $28 \pm 0.5^\circ \text{C}$ and 60-70 % level respectively and food was provided as per OECD guidelines (2004). Hatchlings per box were counted on the 56th day to determine the number of juveniles.

The final endpoints studied were change in biomass on the 28th day, cocoons production per week and number of juveniles on the 56th day. Adult earthworm specimens removed from the test boxes on the 28th day of the experiment were also used to determine the activities of the alkaline phosphatase and total protein of the earthworms following the methods of Walter and Schutt, 1974 and Lowry et al., 1951 respectively. The data was analyzed for single factor ANOVA followed by Least Significance Difference (LSD) test to determine significant variation between treatments at 5 % of probability (Gomez and Gomez, 1984).

RESULTS AND DISCUSSION

There was no mortality in any of the worms in control sets and in T₂, T₃. Average mortality recorded in T₄ was 3.33 %. The control worms showed an increase of 68.1 ± 6.22 % of biomass from the initial value during the 4-weeks test period. The worms exposed to T₂-T₄ doses of the insecticides did not record significant alteration in change of biomass as compared to control. Worms recorded 63.2 ± 3.09 increase in biomass when exposed to the lowest dose, i.e. T₂ representing 12.5% of LC_{50} whereas an increase of 59.19 ± 4.89 % increase of biomass of the test worms was observed when exposed to T₃ dose representing 25% of LC_{50} . Earthworms exposed to T₄ dose of cypermethrin (50% of LC_{50}) recorded 59.21 ± 5.26 % increases in biomass of the test worms from their initial value. This increase was again not significantly different from the change in biomass of the control worms (Fig 1).




Fig. 1.

Number of cocoons produced per worm per week and juveniles produced in control and different treatments on *E. fetida* have been presented in Table 2.

Effects on reproductive parameters

Table 2. Cocoon production and juvenile production in *E. fetida* exposed to control (T1) and sub-lethal doses (T2, T3 & T4) of cypermethrin

Treatment	Cocoons/worm/week*	Juveniles (No. per ten worms)*
T1	1.04 ± 0.05^a	31.0 ± 1.58^a
T2	1.02 ± 0.04^a	30.80 ± 2.40^a
T3	0.99 ± 0.08^a	27.40 ± 0.01^a
T4	0.47 ± 0.01^b	19.80 ± 3.10^b

*Values are expressed as mean \pm S.D; Superscripts in the same column denote significant difference at $p < 0.05$

Mean number of cocoons produced per worm per week in the control was 1.04 ± 0.05 . Mean number of juveniles produced in control was 31.0 ± 1.58 . Sub-lethal doses of the insecticides did not significantly affect the reproductive performance of the worms except at the highest dose (T4) of the insecticide (0.47 ± 0.04 , cocoons/worm/week and juvenile production decreased to 19.80 ± 3.1). Activities of the enzyme alkaline phosphatase of *E. fetida* were significantly elevated in all the treatments except in T2 where the increase was statistically insignificant in comparison to control (Fig 2).




Fig. 2.

Ingesfield (1984) reported low toxicity of cypermethrin to earthworms and explained that this non toxicity can be due to the property of pyrethroids being adsorped onto the organic matter of the soil particles which renders part of the dose unavailable to the worms. Since the LC_{50} value was many times greater than the maximum recommended agricultural dose for the compound, it was concluded by Ingestfield (1984) that cypermethrin would not have adverse effects on natural populations of earthworms. Similar conclusion was made by Dasgupta *et al.*, 2010. Paula *et al.*, 2011 found that earthworms *Eisenia andrei* avoided the soils treated with cypermethrin at concentration $15 \mu\text{g}$ soil onwards but there was no dose-related response within the range of concentration tested. Growth has been reported to be a sensitive parameter to evaluate the toxicity of insecticides on earthworms (Xiao *et al.* 2006). The weight loss may indicate feeding inhibition, with the earthworms regulating the intake of insecticides by reducing consumption rate and thus affecting growth rate (Mosleh *et al.*, 2002, 2003).

Reduction in growth of earthworm by sub-lethal doses of insecticides has also been observed in *Eisenia fetida* (Yasmin and Dsouza, 2007) exposed to several carbamate and organophosphate insecticides. However no significant alteration in the change of biomass of the earthworms could be detected in the present study as compared to control even with the exposure of the highest sub-lethal dose used, i.e., 0.25 mg/kg cypermethrin corresponding to 50% of the LC_{50} value of the insecticide indicating its less toxic nature in comparison to carbamates and organophosphates. Significant alterations in the reproductive performance of the test worms in terms of cocoons/worm/week and juvenile production per ten worms compared to control was detected when exposed to the highest sub lethal dose used, i.e., 0.25 mg/kg cypermethrin corresponding to 50% of the LC_{50} value of the insecticide in the present study indicating its possible hazardous nature. Various scientists have also reported similar result regarding

insecticides influencing reproduction of earthworms (Mosleh *et al.*, 2002; Booth *et al.*, 2000; Yasmin and DSouza, 2007). The observed rise in alkaline phosphates activity in the present study may be indicative of an adaptive rise in enzyme activity to the persistent stress (Murphy and Porter 1966). It is established that insecticide exposure causes reduction in soluble protein and there is a relationship between growth rate, protein content and phosphatases and thus an increase in phosphatases may result. Again elevation of alkaline phosphatase has been reported to be related to resistance of the organism towards the insecticide and the level of pathological and physiological damage caused to the particular organism (Gupta *et al.* 1985; Srinivas *et al.* 2003).

Conclusion

Thus it can be concluded from the present study that sub-lethal dose of cypermethrin significantly affects reproductive performance and alkaline phosphatase activity of the test specimen and their widespread application in agro-ecosystems must be carefully monitored. The present study also indicated the potential use of alkaline phosphatase as a biomarker in assessment of insecticide pollution.

Acknowledgement

We thank the Teacher in Charge, Raja N.L. Khan Women's College for providing necessary laboratory facilities. Financial assistance from University Grants Commission, New Delhi is thankfully acknowledged.

REFERENCES

- Booth, L.H., Hodge, S., O'Halloran, K. 1994. Use of cholinesterase in *Aporrectodea caliginosa* (Oligochaeta; Lumbricidae) to detect organophosphate contamination: Comparison of laboratory tests, mesocosms, and field studies. *Environ Toxicol Chem.*, 19: 417-422.
- Cox, C. 1996. Insecticide Factsheet: Cypermethrin. *J. Pest. Ref.*, 16: 15-20.
- Dasgupta, R., Chakravorty, P. P., Kaviraj, A. 2010. Studies on Relative Toxicities of Six Insecticides on Epigeic Earthworm, *Perionyx excavatus*. *Bull. Environ. Contam. Toxicol.* 85: 83-86.
- Gomez, K. A. and Gomez, A. A. 1984. Statistical procedures for agricultural research; 2nd ed., Wiley-Interscience: New York.
- Gupta, S. K. and Saxena, P. N. 2003. Carbaryl induced behavioral and reproductive abnormalities in the earthworm *Metaphire posthuma*: a sensitive model. *Altern. Lab. Anim.* 31(6): 587-593.
- Ingesfield, C. 1984. Toxicity of the Pyrethroid Insecticides Cypermethrin and WL85871 to the Earthworm, *Eisenia foetida* Savigny. *Bull. Environ. Contam. Toxicol.* 33: 568-570.
- Lowry, O. H., Roberts, N. R., Mei-Ring, W. U., Hixon, W. S., Crawford, E. J. 1951. The quantitative histochemistry of brain enzyme measurements. *J. Bio. Chem.* 207: 19.
- Mosleh, Y. Y., Paris-Palacios, S., Couderchet, M., Vernet, G. 2003 a. Acute and sub-lethal effects of two insecticides on earthworms (*Lumbricus terrestris*) under laboratory conditions. *Environ. Toxicol.* 18(1): 1-8.

- Mosleh, Y. Y., Paris-Palacios, S., Couderchet, M., Vernet, G. 2002. Biological effects of two insecticides on earthworms (*Lumbricus terrestris* L.) under laboratory conditions. Mededelingen Rijksuniversiteit te Gent. Fakulteit van de Landbouwkundige en Toegepaste Biologische Wetenschappen. 67(2): 59–68.
- Mosleh, Y. Y., Paris-Palacios, S., Couderchet, M., Vernet, G. 2003 b. Effects of the herbicide isoproturon on survival, growth rate, and protein content of mature earthworm (*Lumbricus terrestris* L.) Under laboratory conditions. *Enviro. Toxicol.* 18: 1-8.
- Murphy, S. D. and Porter, S. 1966. Effects of toxic chemicals on some adaptive liver enzymes, liver glycogen and blood glucose in fuster rats. *Biochem. Pharmacol.* 15: 1655-1676.
- Nath, B. S., Suresh, A., Varma, B. M., Kumar, R. P. S. 1997. Changes in protein metabolism in hemolymph and fat body of the silkworm, *Bombyx mori* (Lepidoptera: Bombycidae) in response to organophosphorous insecticides toxicity. *Ecotoxicol. Environ. Saf.* 36: 169-173.
- Nowak, J. T., Mccravy, K. W., Fettig, C. J., Berisford, C. W. 2001. Susceptibility of adult hymenopteran parasitoids of the nantucket pine tip moth (Lepidoptera: Tortricidae) to Noctuidae). *J. Econ. Entomol.* 94: 868-873.
- OECD, 1984. Chemicals No 207, Earthworm Acute Toxicity Test. OECD, Paris, France.
- OECD, 2004. Guidelines for testing of chemicals No 222, Earthworm Reproduction Test (*Eisenia fetida/andrei*). OECD, Paris, France.
- Paula, A., Sousa, D., Mara, M., Andréa, D. 2011. Earthworm (*Eisenia andrei*) Avoidance of Soils Treated with Cypermethrin Sensors. 11: 11056-11063.
- Reinecke, S.A. and Reinecke, A. J. 2007. The impact of organophosphate pesticides in orchards on earthworms in the Western Cape, South Africa. *Ecotoxicol. Environ. Saf.* 66: 244-251.
- Srinivas, R., Udikari, S. S., Jayalakshmi, S. K., Sreeramulu, K. 2003. Identification of Factors Responcible for Insecticid Resistance in *Helicoverpa armigera* (Hubner). *Resist. Pest Manag. New.* 13(1): 59-64.
- Suh, C. P. C., Orr, D. B., Van Duyn, J. W. 2000. Effect of insecticides on *Trichogramma exiguum* (Hymenoptera; Trichogrammatidae) preimaginal development and adult survival. *J. Econ. Entomol.* 94: 1122-1129.
- Takada, Y., Kawamura, S., Tanaka, T. 2001. Effects of various insecticides on the development of the egg parasitoid *Trichogramma dendrolimi* (Hymenoptera: Trichogrammatidae). *J. Econ. Entomol.* 94(6): 1340-3.
- Tillman, P. G. and Mulrooney, J. E. 2000. Effect of selected insecticides on the natural enemies *Coleomegilla maculate* and *Hippodamia convergens* (Coleoptera: Coccinellidae), *Geocoris punctipes* (Hemiptera: Lygaeidae), and *Braconmellitor*, *Cardiochiles nigriceps*, and *Cotesia marginiventris*(Hymenoptera : Braconidae) in cotton. *J. Econ. Entomol.* 93: 1638-1643.
- Usmani, K. A. and Knowles, C. O. 2001. Toxicity of pyrethroids and effect of synergists to larval and adult *Helicoverpa zea*, *Spodoptera frugiperda*, and *Agrotis ipsilon* (Lepidoptera: Noctuidae). *J. Econ. Entomol.* 94(4): 868-73.
- Vijenberg, H. P. M. and Van den Bercken, J. 1990. Neurotoxicological effects and the mode of action of pyrethroid insecticides. *Crit. Rev. Toxicol.* 21: 105-126.
- Walkey, A. and Black, I. 1934. An examination of Degtjareff method for determining organic carbon in soils: effect of variations in digestion conditions and of inorganic soil constituents. *Soil Sci.* 63: 251-263.
- Walter, K. and Schutt, C. 1974. Acid and alkaline phosphatase in serum In: Bergmeyer HU (Ed) Methods of enzymatic analysis: New York: Academic Press Inc, pp. 856-864
- Willrich, M. M. and Boethel, D. J. 2001. Effects of diflubenzuron on *Pseudoplusia includens* (Lepidoptera: Noctuidae) and its parasitoid *Copidosoma floridanum* (Hymenoptera: Encyrtidae). *Environmental Entomology.* 30(4): 794-797.
- Xiao, N., Jing, B., Ge, F., Liu, X. 2006. The fate of herbicide acetochlor and its toxicity to *Eisenia fetida* under laboratory conditions. *Chemosphere.* 62(8): 1366–1373.
- Xu, J., Shelton, A. M., Cheng, X. 2001. Variation in susceptibility of *Diadegma insulare* (Hymenoptera: Ichneumonidae) to permethrin. *J. Econ. Entomol.* 94: 541-546.
- Yasmin, S. and DSouza, D. 2007. Effect of Pesticides on the reproductive output of *Eisenia fetida*, *Bull. Environ. Contam. Toxicol.* 79: 529-532.
- Yasmin, S. and DSouza, D. 2010. Effects of pesticides on the growth and reproduction of earthworm: A Rev. *Appl. Environ. Soil Sci.*
- Zhu, G., Wu, H., Guo, J., Kimaro, F. M. E. 2004. Microbial degradation of fipronil in clay loam soil. *Water Air Soil Pollut.* 153(1): 35-44.
